



Jun Haeng Lee <stomachlee@gmail.com>

Description exercise 1

1 개의 메일

Jun Haeng Lee <stomachlee@gmail.com>

2017년 7월 13일 오후 2:05

받는사람: 이준행 Lee Jun Haeng <stomachlee@gmail.com>

하나는 길고 하나는 짧습니다.
1.5cm in length

Case 1

a. 2 linear longitudinal erosions (mucosal breaks) around 1.5cm in length on distal esophagus

b. The 2 longitudinal erosions have edematous mucosal margins with pale discoloration. There is no convergence of folds.

c. Impression: Reflux esophagitis

d. Classification: LA grade B, Modified S-M 1st degree

Modified Savary Miller 분류
쓰신 것 좋기는 한데요...
모름은 거의 안 쓰입니다.
그냥 LA 분류만 쓰면
충분해요

Case 2

a. ulcerated mass around 1.5x3cm along the mid esophagus.

b. Well circumscribed mass with a central deep ulcer that has whitish exudate proximally and a brown exudate distally. The margin is clear and edge is irregular. There is no convergence of folds.

c. Advanced esophageal cancer

d. Bormann type II

매우 좋은데요...

위는 보만 type 을 식도에서 쓰기는 한데요...
약간 어색하다고 생각하는 분이 많아서 그냥
type II 라고 하면 충분

Case 3

a. Multiple converging nodular masses with multiple sizes covering the gastric antrum and the pre-pyloric region.

b. Overall, the region seems edematous with slight hyperemic change in coloration. The nodular masses have smooth surfaces but irregular contour. Many are converging creating merged elevations. The pre-pyloric region is relatively less affected than the distal antrum.

c. Impression: Early gastric cancer

d. Bormann type I + IIa

Mass 는 관리적으로 상당히 큰 흑을
말하므로 ASCO 에서 쓰는 term 인데요.
이것도 같은 이유로
그냥 "ulcer" 라고만 하세요.

Metaplastic gastritis

Case 4

a. 2cm sized ulcerated mass with white exudate in the center of the gastric angle.

b. The margin is irregular, with slight hyperemic and hyperplastic dysplasia with smooth surface. The edge is relatively sharp. There is a slight hyperemic change in the surrounding mucosa.



- c. Impression: Benign gastric ulcer
- d. Classification: Active A2

ulcer는 원래 depressed lesion 이므로
 깊고 shallow ulcer 또는 deep
 erosion 이라고 써 보시오.

Case 5

- a. 3x4cm mass with a depressed ulcer of 0.5x0.5cm on lesser curvature of proximal antrum.
- b. The ulcer is covered with pale whitish exudate. The edge of ulcer is relatively sharp but irregular with hyperemic margin. The masses have smooth surfaces and their contours are well defined. There is convergence of folds toward the ulcer.

- c. Impression: Depressed type of early gastric cancer
- d. IIa+IIc EGC

EGC 에서는 피할 용어임.

Case 6

- a. 5x6cm? mass with central depressed ulceration on antrum
- b. The ulcer is hemorrhagic and has an irregular surface. The edge is not sharp but contained in the depression. The elevated margin is irregular with smooth surface.
- c. Impression: Advanced gastric cancer
- d. Bormann type II

Case 7

- a. Duodenal bulb thickening
- b. The mucosa has a pale coloration on the contour closer to lumen, and a hyperemic coloration on the outer mucosa.
- c. Impression: Duodenal ulcer scar

- d. Scar stage 1 and 2

regenerating epithelium 처럼 보이기
 붉은 색이면 S1, 주변과 거의
 비슷하거나 오히려
 pale하면 S2로
 알려주세요.

이준행, 李竣行, Jun Haeng Lee, M. D.
 Department of Medicine, Samsung Medical Center, Sungkyunkwan University
 Address: (ZIP: 06351) 83 Irwon-ro Gangnam-gu, Seoul, Korea
 Telephone: 02-3410-3409, 0912
 Homepage: <http://endotoday.com>

매우 잘했습니다.

2017-7-13. 이준행